

Guidance: treating enteral stoma site overgranulation

v2 May 2022

GP / DOCTOR	<ul style="list-style-type: none">• Refer patient to GP/ ward doctor for initial review of stoma site• Follow guidance as per steps below
TREAT	<ul style="list-style-type: none">• Clean site with Prontosan wash twice daily• Apply steroid cream according to grade of overgranulation (see below)• Ensure this is applied to granulomatous tissue only, not to normal skin• Review stoma site at least every 2 weeks and adjust treatment accordingly
REVIEW	<ul style="list-style-type: none">• Apply Polyurethane Foam dressing (ie Aquacel foam non adhesive 5cm; Allevyn adhesive 7.5cm) as a compressive keyhole dressing twice daily or as needed. If skin sensitive to dressings, use dry gauze as keyhole dressing.• Ensure the fixation plate is secured firmly over the top of the dressing• Ensure the tube itself is secured to the abdomen (eg with Cinch device)
Step 1: mild overgranulation	<ul style="list-style-type: none">• Mild potency steroid cream such as 1-2.5% Hydrocortisone cream applied twice daily after cleaning affected area with Prontosan wash• If no improvement after 2 weeks, treat as for step 2
Step 2: moderate overgranulation	<ul style="list-style-type: none">• Moderate potency steroid cream such as Clobetasone Butyrate (Eumovate) applied twice daily after cleaning affected area with Prontosan wash• Review at least every 2 weeks• If no improvement after 6 weeks, treat as for step 3
Step 3: severe overgranulation	<ul style="list-style-type: none">• Potent steroid cream such as Betamethasone 0.1% applied twice daily after cleaning affected area with Prontosan wash• Review at least every 2 weeks
HEALED	<ul style="list-style-type: none">• Once stoma site has healed ensure prescribed steroid cream and dressings are discontinued and Prontosan wash amended to once daily maintenance dose
MONITOR	<ul style="list-style-type: none">• Continue to monitor the stoma site daily for signs of the overgranulation returning and ensure that the tube is secured firmly to the abdomen at all times
REFER	<ul style="list-style-type: none">• If no improvement to stoma site having treated as for steps 1-3, complete the enteral referral form on EPR